

# Determinants of Ruptured Cerebral Aneurysm and the Presenting Symptoms among Patients with SAH Admitted at a Tertiary Care Center in North India

Mahendra Kumar<sup>1</sup>, Sukhpal Kaur<sup>2</sup>, Ashish Aggarwal<sup>3</sup>, Pravin Salunke<sup>4</sup>, Monaliza Mittal<sup>5</sup>

## ABSTRACT

**Introduction:** Aneurysmal subarachnoid hemorrhage (SAH) is a serious disease with high rates of mortality and morbidity. The current study is carried out to assess the risk factors and clinical profile of the patients with aneurysmal SAH presenting to a tertiary care center in India.

**Materials and methods:** A single-center cross-sectional study was carried out on 65 cerebral aneurysm patients with an aneurysmal SAH. The sociodemographic details, comorbidities, risk factors, and the presenting symptoms experienced by the patients at the time of rupture of aneurysm were recorded by using a pre-validated questionnaire.

**Results:** The age of the participants ranged between 41 and 55 years and the majority were female. Maximum had experienced severe headache followed by vomiting and neck stiffness before rupturing of the aneurysm. Age above 45 years followed by hypertension, female gender, and smoking were identified as major risk factors among the study subjects.

**Conclusion:** Advanced age, smoking, and female gender are found main risk factors with severe headache, vomiting, and neck rigidity as main symptoms. Risk factors and symptoms are important to know for future research and preparing treatment modules.

**Keywords:** Cerebral aneurysm, Risk factors, Ruptured symptoms.

*Pondicherry Journal of Nursing* (2021): 10.5005/jp-journals-10084-12172

## INTRODUCTION

A cerebral aneurysm is one of the major cerebrovascular diseases causing detrimental effects on individuals' health and performance due to rupturing and creating subarachnoid hemorrhage (SAH).<sup>1</sup>

A brain aneurysm is an abnormal focal dilation of a blood vessel wall in the brain resulting from the weakening of the inner muscle layer of the blood vessel wall.

Aneurysm rupture with resultant SAH is the most feared consequence of unruptured intracranial aneurysms. This wall gets weakened by injury or any disease.<sup>1,2</sup>

Aneurysm rupture is one of the most intricate complications of cerebral aneurysms and about 80% of non-traumatic SAHs are related to rupture of cerebral aneurysms where the mortality rate is as high as 45%.<sup>2-4</sup> Prompt detection and timely treatment are censorious for good recovery and outcomes in patients with non-traumatic SAH.<sup>4</sup>

Usually, aneurysms remain silent till rupture and then present with acute and variety of symptoms including sudden headache and neck rigidity. Focal neurological deficits are likely due to an increase in mass effect, or any event of cerebral ischemia distal to the site of the aneurysm.<sup>5</sup>

Patients who present to triage with symptoms, such as, acute loss of consciousness, severe headache which is often accompanied by vomiting, and neck rigidity indicate the presence of aneurysmal SAH.<sup>5</sup> In 30% of ruptured cerebral aneurysm patients, headache is usually lateralized to the side of the aneurysm.<sup>6,7</sup>

Subarachnoid hemorrhages most often with cerebral aneurysms are silent and present with a bleed in the brain.<sup>8</sup> A study conducted in Japan reported >90% of cerebral aneurysm cases detected incidentally raise the need for an institutional systemic

<sup>1</sup>Department of Neurology, Postgraduate Institute of Medical Education and Research, Chandigarh, India

<sup>2,5</sup>Department of Nursing, Postgraduate Institute of Medical Education and Research, Chandigarh, India

<sup>3</sup>Department of Neurosurgery, Postgraduate Institute of Medical Education and Research, Chandigarh, India

<sup>4</sup>National Institute of Nursing Education, PGIMER, Chandigarh, India

**Corresponding Author:** Mahendra Kumar, Department of Neurology, Postgraduate Institute of Medical Education and Research, Chandigarh, India, Phone: +91 9465223015, e-mail: kumarmahen9@gmail.com

**How to cite this article:** Kumar M, Kaur S, Aggarwal A, Salunke P, Mittal M. Determinants of Ruptured Cerebral Aneurysm and the Presenting Symptoms among Patients with SAH Admitted at a Tertiary Care Center in North India. *Pon J Nurs* 2021;14(1):2-7.

**Source of support:** Nil

**Conflict of interest:** None

approach of preventive methods, such as, mass screening to detect unruptured aneurysms.<sup>9</sup>

A raised incidence of the ruptured cerebral aneurysm has been found in patients with certain predisposing factors, such as, advanced age, hypertension, smoking, alcohol abuse, and atherosclerosis.<sup>2,10</sup> Cigarette smoking and hypertension are among the leading influencing factors of non-traumatic SAH.<sup>11,12</sup>

The peak incidence of cerebral aneurysm rupture is among the age group between 40 and 60 years, with 53 years as the mean age.<sup>11</sup> A meta-analysis on aneurysmal SAH depicts that this disproportionately affects more to women likely causes being stress, reproductive factors, and some other genetic variations expanding the risk in women.<sup>13</sup>

Further aggravating factors include alcohol habits, personal or family history of aneurysm, hemorrhagic stroke, and cerebrovascular disease.<sup>11,14</sup> Diabetic mellitus is another well-known predisposing factor for cerebral infarction.<sup>15-18</sup> World Federation of Neurosurgical Societies (WFNS), Hunt and Hess, and Fisher grade scales are worldwide being used to observe the severity of ruptured aneurysm and give a prediction of treatment outcome.<sup>19</sup>

### AIM OF THE STUDY

The present study aimed to predict the determinants of ruptured cerebral aneurysms among patients presenting at a tertiary care hospital in North India.

### MATERIALS AND METHODS

A single-center cross-sectional study was carried out at a tertiary care center in Chandigarh, India.

Sixty-five ruptured cerebral aneurysm patients presented in the neurosurgery unit of the institute from June 2017 to September 2017 were enrolled in the study. For all the ruptured aneurysm, CT was used for initial diagnosis and treatment decision-making. The identification data of patients with ruptured cerebral aneurysms treated/presented at the mentioned center were extracted from the list of admission records. The required information was obtained from the next kin of the patients, an archive of hospital records, and documented in the study questionnaire. Ethical permission was obtained from the Institutional Ethics Committee. The anonymity of the information of the patient was ensured. Written informed consent was taken from all the patients or their next of kin, as applicable.

#### Inclusion Criteria

All ruptured cerebral aneurysm patients presented to the given tertiary care center during the study period.

#### Exclusion Criteria

Subarachnoid hemorrhage due to any other cause (trauma, intracranial arteriovenous malformations, hypertension, and coagulopathy).

#### Questionnaire


A pre-validated questionnaire was used to collect the information from study subjects including demographic details, such as, the age of patients, gender, and occupation. The clinical profile consisted of symptoms experienced by the patients at the time of aneurysm rupture, risk factors, such as, comorbidities, and severity grading of aneurysmal subarachnoid hemorrhage (aSAH) as per WFNS, H and H, and Fisher grade scales. The clinical characteristics analyzed as a known case of hypertension (we kept diagnosed hypertension as time after time systolic blood pressure (SBP) >140 mm Hg or diastolic blood pressure over 90 mm Hg, or on blood pressure control medication), diabetics, smoking habits (ex-smokers or history of smoking over 10 years, and continued smokers), and habit of alcohol intake.

### RESULTS

Table 1 depicts the demographic and clinical characteristics of the patients. The mean ± SD age of the patients was 49.82 ± 11.61 years. The majority of the patients were female (56.9%). 41.5% of study participants were educated up to senior secondary whereas

**Table 1:** Sociodemographic determinants of study participants (N = 65)

Variable	n (%)
Age (years)	
18–29	06 (9.3)
30–44	10 (15.4)
45 years and above	49 (75.3)
Age: mean ± SD: 49.82 ± 11.61	
Range: 18–64 years	
Gender	
Male	28 (43.1)
Female	37 (56.9)
Educational status	
Just literate	20 (30.8)
Up to middle class	18 (27.7)
Senior secondary and above	27 (41.5)
Area of living	
Rural	55 (84.6)
Urban	10 (15.4)
Religion	
Hindu	51 (78.5)
Sikh	8 (12.3)
Others (Muslim, Christian)	06 (9.2)
Type of family	
Nuclear	11 (16.9)
Joint	54 (83.1)
Occupation	
Not working/homemaker	40 (61.5)
Skilled/unskilled worker	06 (9.2)
Govt. employee	09 (13.8)
Agriculture	10 (15.4)



**Fig. 1:** Symptoms of ruptured cerebral aneurysm (N = 65)

30.8% were just literate. Maximum (84.6%) of participants were from rural areas and were Hindu (78.5%) by religion. 61.5% of the patients were either not working or homemakers as more than half of the patients were female.

Figure 1 shows that 95.4% of the patients experienced a headache after the rupture of aneurysm followed by vomiting (72.3%). Neck stiffness was reported by 64.6 and 46.4% of

participants reported irritability. Loss of consciousness was reported among 44.6% of patients with a mean  $\pm$  SD duration of  $4.82 \pm 6.92$ .

Hypertension was reported by 64.6% with the mean  $\pm$  SD duration of  $4.35 \pm 3.79$  years of the disease. History of diabetes mellitus was reported by 9.2% with the mean  $\pm$  SD duration of  $7.67 \pm 2.73$  years. Treatment adherence was better in the case of diabetes mellitus (83%) as compared to hypertension. All the patients suffering from seizures and chronic obstructive pulmonary disease (COPD) had good adherence to treatment (Table 2).

Advanced age, above 45 years was the major risk factor reported from study results as 75.3% of patients were above 45 years of age and the second main risk factor was the female gender (56.9%). The habit of smoking was reported by 29.2% with the mean  $\pm$  SD duration of  $3.20 \pm 6.16$  years. 21.5% had a history of alcohol consumption (Figs 2 and 3).


Table 3 depicts the severity of ruptured cerebral aneurysm as assessed by three scales. As per Hunt and Huss scale, 58.7% were in grade I with the mean  $\pm$  SD score of  $2.29 \pm 0.669$ . World Federation of Neurological Societies grade I was 38.5% with a mean  $\pm$  SD score of  $1.89 \pm 0.868$  as leading. Equally, 41.7% of patients had Fisher grade I, grade III SAH on CT scan, and mean  $\pm$  SD was  $3.20 \pm 0.887$ . A total of 56.92 patients were having an aneurysm at the anterior cerebral artery/internal carotid artery (ACA/ICA) location.

Out of 65 patients, 36 (55.4%) were conscious at presentation. The mean duration  $\pm$  SD of loss of consciousness (LOC) was  $4.82 \pm 6.92$ , whereas 55.4% did not report LOC. 18.4% of study participants had 6 to 10 minutes duration of LOC. The majority of the aneurysm (56.9%) were located at ICA/ACA and the maximum (87.7%) were secular aneurysms (Table 4).

78.5% of subjects reached the tertiary healthcare centers on the same day and 10.7% reached within 5–7 days with a mean  $\pm$  SD of  $2.74 \pm 1.98$ .

**Table 2:** Risk factors (comorbidities) associated with ruptured cerebral aneurysm (N = 6)

Variables	N (%)	Duration comorbidities (Mean $\pm$ SD)	Treatment adherence
Hypertension	42 (64.6)	$4.35 \pm 3.79$	24 (57.14%)
Diabetic mellitus	06 (9.2)	$7.67 \pm 2.73$	05 (83%)
Seizure	02 (3.1)	$2.20 \pm 3.19$	02 (100%)
COPD	02 (3.1)	$12.50 \pm 3.53$	02 (100%)



**Fig. 2:** Risk factors of ruptured cerebral aneurysm (N = 65)

## DISCUSSION


Subarachnoid hemorrhage is among the most destructive neurological diseases. This disease not only impacts the consequential effect on morbidity and mortality but also conveys awful social and financial consequences.<sup>3</sup>

The present study was carried out at a well-known over 2,000 beds capacity tertiary health care center in Chandigarh, India. It is also known as a center for excellence with mass volume in neurosurgical cases. Aneurysmal SAH is a catastrophic acute neurological disease that impacts over 30,000 population annually in the USA.<sup>4</sup>

The intrinsic complexities of the treatment have a high impact on advancements in investigation facilities and treatment modalities mean that these patients can be more effectively treated at tertiary care centers. According to the results obtained from our study and other studies, age is a factor affecting the incidence of an aneurysm. About two-thirds (75.3%) of study participants were above the age of 45 years with an average age of  $49.82 \pm 11.61$

**Table 3:** Clinical severity grading of ruptured cerebral aneurysm (N = 65)

Clinical grading	n (%)
Hunt and Huss scale	
Grade I	05 (7.7)
Grade II	38 (58.7)
Grade III	20 (30.8)
Grade IV	02 (3.1)
Mean $\pm$ SD	$2.29 \pm 0.669$
WFNS grade (at presentation)	
Grade I	25 (38.5)
Grade II	24 (36.9)
Grade III	15 (23.1)
Grade IV	01 (1.5)
Mean $\pm$ SD	$1.89 \pm 0.868$
Fisher: CT grade	
Grade I	27 (41.7)
Grade II	07 (10.8)
Grade III	27 (41.7)
Grade IV	04 (6.1)
Mean $\pm$ SD	$3.20 \pm 0.887$



**Fig. 3:** Time duration to reach tertiary health care (N = 65)

**Table 4:** Other variables of ruptured aneurysm patients  $N = 65$ 

	Variable	N (%)
1	Duration of LOC	
	No LOC	36 (55.4)
	1–5 minutes	10 (15.3)
	6–10 minutes	12 (18.4)
	11 minutes or above	07 (10.7)
	Mean $\pm$ SD	4.82 $\pm$ 6.92
2	Location of aneurysm	
	ICA/ACA	37 (56.92)
	A.com.	14 (21.53)
	MCA	09 (13.8)
	Posterior circulation/others	05 (7.6)
3	Type of aneurysm	
	Secular	57 (87.7)
	Fusiform	02 (3.1)
	Others (specify)	06 (9.2)
4	Time duration to reach tertiary health care	
	0–1 day	51 (78.5)
	2–4 days	08 (12.3)
	5–7 days	07 (10.76)
	Mean $\pm$ SD	2.74 $\pm$ 1.98

years. Aneurysmal SAH has a peak incidence of 40 to 60 years of age.<sup>11</sup> According to Cowan et al., the mean age of patients with an aneurysm was 53.2 years,<sup>20</sup> which is almost similar to that of our study findings. One of the largest studies, regarding cerebral aneurysm, reported that advanced age is the most significant risk factor associated with surgical outcome.<sup>21</sup>

Gender has a marked dominancy on the incidence of the cerebral aneurysm as they that found women have 1.6 times higher risk than men and prevalence of cerebral aneurysm in male and female was reported as 2:1.<sup>22</sup> Female gender was dominating as 56.9% of the total study population were females in our study. Another study has also revealed that the female gender represents a significant risk factor as women outnumbered men among aSAH patients, especially along with increasing age strata.<sup>23</sup>

Headache in an acute and severe degree at the time of Ictus was the most common symptom registered as 95.4% of patients experienced a headache. Similar findings were reported by another study as the classic symptoms of aneurysmal rupture characterized by a sudden and severe headache which turn in the majority of cases at its utmost severity with no time and is often reported by patients as “the worst headache of my life”, may last for a few hours or days.<sup>6,17</sup> Both, the intensity and acute onset of severe headache are highly characteristic features of a ruptured cerebral aneurysm. Patients who already suffered from chronic headache before aneurysm rupture usually reported pain with an utmost difference in nature and intensity.<sup>24</sup>

Loss of consciousness is among the most common presenting symptoms of non-traumatic SAH.<sup>25</sup> In our study, 44.6% of participants had experienced the loss of consciousness with the mean  $\pm$  SD duration of 4.82  $\pm$  6.962. Other risk factors including seizure and COPD were almost equally distributed as 3.1% of patients.

A similar study conducted by Longstrn and colleagues supported our study findings as alcohol consumption, cigarette

smoking, and hypertension have been studied as leading independent risk factors for SAH.<sup>22</sup> Our study results show that 21.5% of patients were having a history of alcohol intake with 1.43  $\pm$  3.321 of mean  $\pm$  SD duration. A long duration of heavy alcohol consumption is significantly associated with cerebral aneurysm rupture and the leading cause of transient increases in blood pressure.<sup>26,27</sup>

Studies have found smoking as the most important established risk factors for aneurysmal SAH.<sup>6,12</sup> Our study reported 29.5% of participants had a history of smoking with the mean  $\pm$  SD 3.20  $\pm$  6.162 duration. Cigarette smoke may lead to cerebral aneurysm genesis and rupture through several mechanisms as may influence each step in the cascade of events leading to aneurysm evolution, growth, and rupture recently reviewed.<sup>27</sup> Patients harboring cerebral aneurysm should be strongly counseled to quit smoking and alcohol intake. The importance of this simple intervention cannot be overemphasized and counseling at the time of discharge and OPD visits.

Systemic hypertension has long been considered a primary risk factor of aneurysmal rupture. Hypertension was one of the most reported modifiable risk factors (64.6%) found in our study and treatment adherence was 57.14% that shows a lack of treatment continuity among people suffering from hypertension.<sup>3,28</sup> Most of the patients with a known history of hypertension are usually put on antihypertensive agents. So, they tend to have a normal range of blood pressure reading before an aneurysm diagnosed.<sup>21</sup>

Hence, blood pressure control is a pivotal practice that helps to control the risk of aneurysm rupture and medication adherence is needed to maintain as hypertension had been a positive effect on suffering to cerebral aneurysms.

Expanding population, ever-changing way of life, population aging, diabetes mellitus is the most common metabolic syndrome and a prime source of disability worldwide. In our study, 9.2% of patients had diabetes as a comorbid risk.<sup>29</sup> Diabetes impacts the arterial wall and is a well-known trigger for cerebral infarction but that his connection with the development of SAH, as the cause of ruptured aneurysms still remains unclear.<sup>29–31</sup>

Our study participants were from different age groups, backgrounds, and family types. The majority of the patients were residents of the rural area but the effect of geographical region on incidence and type of aneurysm rupture is indecisive at present.<sup>9</sup> 78.5% of patients reached the tertiary care center within 24 hours of ictus.

Assessment of the patient’s severity of condition immediately after a SAH is of the utmost importance and crucial to further management. 56.92% of patients had an aneurysm location at ACA/ICA. Classification of severity of rupture should be done in a reliable way and with relevance to prognosis and outcome. Severe Hunt and Hess grades II was 58.7% generally consist of approximately 20–30% of those admitted to the hospital with aneurysmal SAH 36.9%. The WFNS grading system uses the Glasgow Coma Scale and the presence of focal neurological deficits to grade the severity of SAH. Where assessing patients with a proven aneurysmal SAH, not only the clinical grading is important but also the amount of subarachnoid blood assessed by Fisher scale.

In an attempt to get a sense of the fate that befalls the patients of a ruptured cerebral aneurysm in our country, we studied the contributory factors and clinical profile of patients presenting to a high-volume center for aneurysmal surgery in North India. However, developing countries like India still is lagging in establishing such facilities for their vast population.



Data were collected over 3 months and descriptive analysis was done. It is crucial to acknowledge risk factors related to cerebral aneurysms so that the information can be used to educate to avoid the modifiable risk factors.

## CONCLUSION

Minimizing the risk of formation and rupture of cerebral aneurysm via modifiable risk factors identification and timely management, rather than surgical intervention is of great clinical and social value. Aneurysmal SAH is a life-threatening condition and the present study puts important information regarding symptoms and risk predictors of a ruptured cerebral aneurysm.

## STRENGTH OF OUR STUDY

The scope of studying sociodemographic and clinical profile is not limited to hospital services, and this information can also be utilized while making public health policies and implementation of cerebral aneurysm awareness programs.

## ANALYSIS AND INTERPRETATION

Collected data were analyzed by using descriptive analysis which is presented in the form of tables and figures.

## REFERENCES

- King JT, Horowitz MB, Kassam AB, Yonas H, Roberts MS. The short form-12 and the measurement of health status in patients with cerebral aneurysms: performance, validity, and reliability. *J Neurosurg* 2005;3(3):489–493. DOI: 10.3171/jns.2005.102.3.0489.
- Jill Novitzke RN. The basics of brain aneurysms: a guide for patients. *J Vasc Interv Neurol* 2008. 89–90.
- Bederson JB, Connolly Jr ES, Hunt Batjer H, Dacey RG, Dion JE, Diringer MN, et al. Guidelines for the management of aneurysmal subarachnoid hemorrhage: a statement for healthcare professionals from a special writing group of the stroke council American heart association. *Stroke* 2009;40(3):994–1025. DOI: 10.1161/STROKEAHA.108.191395.
- Rincon F, Rossenwasser RH, Dumont A. The epidemiology of admissions of nontraumatic subarachnoid hemorrhage in the United States. *Neurosurgery* 2013;3(2):217–222. DOI: 10.1227/01.neu.0000430290.93304.33.
- Cohen-Gadol AA, Bohnstedt BN. Recognition and evaluation of nontraumatic subarachnoid hemorrhage and ruptured cerebral aneurysm. *Am Fam Physician* 2013;88:451–456.
- Ju YE, Schwedt TJ. Abrupt-onset severe headaches. *Semin Neurol* 2010;30(02):192–200. DOI: 10.1055/s-0030-1249229.
- Liebenberg WA, Worth R, Firth GB, Olney J, Norris JS. Aneurysmal subarachnoid haemorrhage: guidance in making the correct diagnosis. *Postgrad Med J* 2005;81(957):470–473. DOI: 10.1136/pgmj.2004.023788.
- Pierot LG, Gawlitza M, Soize S. Unruptured intracranial aneurysms: it is not a bomb. *Rev Neurol* 2017;173(9):530–531. DOI: 10.1016/j.neurol.2017.05.018.
- Morita A, Kirino T, Hashi K, Aoki N, Fukuhara S, Hashimoto N, et al., UCAS Japan##The natural course of unruptured cerebral aneurysms in a Japanese cohort. *N Eng J Med* 2012;366(26):2474–2482. DOI: 10.1056/NEJMoa1113260.
- Juvela S, Korja M. Intracranial aneurysm parameters for predicting a future subarachnoid hemorrhage: a long-term follow-up study. *Neurosurgery* 2017;81(3):432–440. DOI: 10.1093/neuros/nyw049.
- Broderick JP, Viscoli CM, Brott T, Kernan WN, Brass LM, Feldmann E, et al. Major risk factors for aneurysmal subarachnoid hemorrhage in the young are modifiable. *Stroke* 2003;34(6):1375–1381. DOI: 10.1161/01.STR.0000074572.91827.F4.
- Kleinloog RD, de Mul N, Verweij BH, Post JA, Rinkel GJE, Ruigrok YM. Risk factors for intracranial aneurysm rupture: a systematic review. *Neurosurgery* 2018;82(4):431–440. DOI: 10.1093/neuros/nyx238.
- Rehman S, Sahle BW, Chandra RV, Dwyer M, Thrift AG, Callisaya M, et al. Sex differences in risk factors for aneurysmal subarachnoid haemorrhage. *J Neurolog Sci* 2019;406:116446. DOI: 10.1016/j.jns.2019.116446.
- Feigin VL, Rinkel GJE, Lawes CMM, Algra A, Bennett DA, van Gijn J, et al. Risk factors for subarachnoid hemorrhage: an updated systematic review of epidemiological studies. *Stroke* 2005;36(12):2773–2780. DOI: 10.1161/01.STR.0000190838.02954.e8.
- Almdal TS, Scharling H, Jensen JS, Vestergaard H. The independent effect of type 2 diabetes mellitus on ischemic heart disease, stroke, and death: a population-based study of 13,000 men and women with 20 years of follow-up. *Arch Intern Med* 2004;164(13):1422–1426. DOI: 10.1001/archinte.164.13.1422.
- O'Donnell MJ, Xavier D, Liu L, Zhang H, Chin SL, Rao-Melacini P, et al. Risk factors for ischaemic and intracerebral haemorrhagic stroke in 22 countries (the INTERSTROKE study): a case-control study. *Lancet* 2010;376(9735):112–123. DOI: 10.1016/S0140-6736(10)60834-3.
- Guimond JG, Chagnon P-M, Bojanowski MW. Clipping vs. Coiling in acute aneurysmal subarachnoid haemorrhage: should the patient's medical condition influence treatment modality. *Neuro-Chirurgie* 2012;58(2-3):115–124. DOI: 10.1016/j.neuchi.2012.02.012.
- de Rooij NK, Linn FHH, van der Plas JA, Algra A, Rinkel GJE. Incidence of subarachnoid haemorrhage: a systematic review with emphasis on region, age, gender and time trends. *J Neurol Neurosurg Psychiatry* 2007;78(12):1365–1372. DOI: 10.1136/jnnp.2007.117655.
- Aggarwal AS, Salunke P, Singh H, Kumar Gupta S, Chhabra R, Singla N, et al. Vasospasm following aneurysmal subarachnoid hemorrhage: thrombocytopenia a marker. *J Neurosci Rural Pract* 2013;4(3):257–261. DOI: 10.4103/0976-3147.118762.
- Cowan J, Dimick JB, Wainess RM, Upchurch Jr GR, Thompson BG. Outcomes after cerebral aneurysm clip occlusion in the United States: the need for evidence-based hospital referral. *J Neurosurg* 2004;99(6):947–952. DOI: 10.3171/jns.2003.99.6.0947.
- Wiebers DO, Whisnant JP, Huston 3rd J, Meissner I, Brown Jr RD, Piepgras DG, et al. Unruptured intracranial aneurysms: natural history, clinical outcome, and risks of surgical and endovascular treatment. *Lancet* 2003;362(9378):103–110. DOI: 10.1016/S0140-6736(03)13860-3.
- Eden SV, Meurer WJ, Sánchez BN, Lisabeth LD, Smith MA, Brown DL. Gender and ethnic differences in subarachnoid hemorrhage. *Neurology* 2008;71(10):731–735. DOI: 10.1212/01.wnl.0000319690.82357.
- De Marchis GM, Schaad C, Fung C, Beck J, Gralla J, Takala J. Gender-related differences in aneurysmal subarachnoid hemorrhage: a hospital based study. *Clin Neurol Neurosurg* 2017;157:82–87. DOI: 10.1016/j.clineuro.2017.04.009.
- Cebral JR, Mut F, Weir J, Putman CM. Association of hemodynamic characteristics and cerebral aneurysm rupture. *AJNR Am J Neuroradiol* 2011;32(2):264–270. DOI: 10.3174/ajnr.A2274.
- Suwatcharangkoon SM, Meyers E, Falo C, Schmidt JM, Agarwal S, Claassen J, et al. Loss of consciousness at onset of subarachnoid hemorrhage as an important marker of early brain injury. *JAMA Neurol* 2016;73(1):28–35. DOI: 10.1001/jamaneurol.2015.3188.
- Can AC, Castro VM, Ozdemir YH, Dagen S, Dligach D, Finan S, et al. Alcohol consumption and aneurysmal subarachnoid hemorrhage. *Transl Stroke Res* 2018;9(1):13–19. DOI: 10.1007/s12975-017-0557-z.
- Chalouhi NA, Chalouhi N, Ali MS, Starke RM, Jabbour PM, Tjoumakaris SI, et al. Cigarette smoke and inflammation: role in cerebral aneurysm formation and rupture. *Mediators Inflamm* 2012;2012:271582. DOI: 10.1155/2012/271582.

28. Nuki YT, Tsou T-L, Kurihara C, Kanematsu M, Kanematsu Y, Hashimoto T. Elastase-induced intracranial aneurysms in hypertensive mice. *Hypertension* 2009;54(6):1337–1344. DOI: 10.1161/HYPERTENSIONAHA.109.138297.
29. Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Res Clin Pract* 2010;87(1):4–14. DOI: 10.1016/j.diabres.2009.10.007.
30. Murray CJL, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010. *Lancet* 2012;380(9859):2197–2223. DOI: 10.1016/S0140-6736(12)61689-4.
31. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2010;33(Suppl 1):S62–S69. DOI: 10.2337/dc11-S062.